

A Comparison
of
Two Models in Physics:
(new and currently accepted)

Dynamic (DM)
and
Standard (SM)

Advantages of the Dynamic Model (DM)

described by classical methods of wave physics, in particular,
by the general wave equation

$$\Delta \hat{\Psi} - \frac{1}{c^2} \frac{\partial^2 \hat{\Psi}}{\partial t^2} = 0$$

What does it follow from the DM?

Comments on capability of the Standard Model (SM)

based on:

Schrödinger's QM, Dirac's QED, and QCD,
with respect to enumerated points at issue

1

The origin of mass.

The mass has the field **associated** character:

$$m = \frac{4\pi\epsilon_0 r^3}{1 + k_e^2 r^2}, \quad \epsilon_0 = 1 \text{ g} \cdot \text{cm}^{-3}, \quad k_e = \omega_e / c \text{ (see \# 5)}.$$

The rest mass does not exist.

Unknown

2

The nature of electric charges.

The charge is the rate of mass exchange:

$$Q = dm / dt$$

Unknown

3

The relation between the mass and charge:

$$Q = m\omega_e$$

Unknown

4

The objective central ("electric") charge of an electron:

$$e = 1.70269155 \cdot 10^{-9} \text{ g} \cdot \text{s}^{-1}$$

Electron is **an elementary quantum of the rate of mass exchange.**

Incorrect dimension and value:

$$e = 1.602176462 \cdot 10^{-19} \text{ C (SI)},$$

$$\text{where } 1\text{C} = \frac{c_r}{10} \frac{1}{\sqrt{10^9}} \text{ kg}^{1/2} \text{ m}^{3/2} \text{ s}^{-1},$$

$$c_r = 2.99792458 \cdot 10^{10} \text{ or in a CGSE system}$$

$$e = 4.803204197 \cdot 10^{-10} \text{ CGSE}_q \text{ (g}^{1/2} \cdot \text{cm}^{3/2} \cdot \text{s}^{-1}\text{)}$$

5	<p>The fundamental frequency of the subatomic and atomic levels:</p> $\omega_e = e / m_e = 1.869162559 \cdot 10^{18} \text{ s}^{-1}$	Unknown
6	<p>Static fields do not exist in Nature.</p> <p>“Electrostatic” fields are, actually, exofrequency fields of the fundamental frequency ω_e (see # 5)</p>	Unknown
7	<p>The objective transversal (“magnetic”) charge of an electron on the Bohr orbit:</p> $e_H = \frac{v_0}{c} e$	Unknown
8	<p>The fundamental wave radius:</p> $\tilde{\lambda}_e = c / \omega_e = 1.603886492 \cdot 10^{-8} \text{ cm}$	Unknown
9	<p>The ratio of electron’s orbital magnetic moment, to its orbital moment of momentum,</p> $\mu_{e,orb} = e v_0 r_0 / c ,$ $\hbar = m_e v_0 r_0 :$ $\frac{\mu_{e,orb}}{\hbar} = \frac{e}{m_e c} = \frac{1}{\tilde{\lambda}_e} = k_e$	<p>Incorrect value</p> $\frac{\mu_{e,orb}}{\hbar} = \frac{e}{2m_e c}$
10	<p>The magnetic moment of an electron:</p> $\mu_e = \frac{v_0}{c} e (r_0 + \delta r_0) = -1855.877359 \cdot 10^{-26} \text{ J} \cdot \text{T}^{-1} ;$ <p>$v_0 = 2.187691263 \cdot 10^8 \text{ cm} \cdot \text{s}^{-1}$ is the Bohr speed</p>	<p>Incorrect value</p> $\mu_e = (1 + \alpha_e) \frac{e \hbar}{2m_e c} =$ $= -928.476410(80) \cdot 10^{-26} \text{ J} \cdot \text{T}^{-1}$

11	<p>The proper magnetic moment of an electron (electron “spin” magnetic moment):</p> $\mu_s = \frac{r_e}{z_{p,q}} \sqrt{\frac{2Rh_e}{m_0c}} = -5.50792 \cdot 10^{-29} \text{ J} \cdot \text{T}^{-1}$	<p>Incorrect value</p> $\mu_s = \mu_B = \frac{e\hbar}{2m_e c} = -927.400947(80) \cdot 10^{-26} \text{ J} \cdot \text{T}^{-1}$
12	<p>The radius of an electron shell (electron’s radius):</p> $r_e = \sqrt{\frac{m_e}{4\pi\epsilon_0}} = 4.17052597 \cdot 10^{-10} \text{ cm};$ $\epsilon_0 = 1 \text{ g} \cdot \text{cm}^{-3}, m_e = 9.10938253 \cdot 10^{-28} \text{ g}$	<p>Unknown</p> <p>Considered as a point like particle. Classical electron radius is</p> $r_e = \left(\frac{v_0}{c}\right)^2 r_0 = 2.817940325 \cdot 10^{-13} \text{ cm}$
13	<p>The radius of a proton shell (proton’s radius):</p> $r_p = 0.528421703 \cdot 10^{-8} \text{ cm}$ <p>(calculated from the formula of mass, see # 1)</p>	<p>Unknown</p> <p>Proton rms charge radius is</p> $r_p = 0.8750(68) \cdot 10^{-13} \text{ cm}$
14	<p>The fundamental frequency of the gravity field:</p> $\omega_g = \sqrt{4\pi\epsilon_0 G} = 9.158082264 \cdot 10^{-4} \text{ s}^{-1};$ $G = 6.6742 \cdot 10^{-8} \text{ g}^{-1} \cdot \text{cm}^3 \cdot \text{s}^{-2}, \quad \epsilon_0 = 1 \text{ g} \cdot \text{cm}^{-3}$	<p>Unknown</p>

15	<p>The fundamental wave radius of the gravity field:</p> $\hat{\lambda}_g = c / \omega_g = 327.4 Mkm$	<p>Unknown</p>
16	<p>The gravitational spectrum of nucleon wave shells:</p> $r = \hat{\lambda}_g z_{m,n};$ <p>$z_{m,n}$ are roots of Bessel functions</p>	<p>Unknown</p>
17	<p>The background spectrum of the hydrogen atom:</p> $\frac{1}{\lambda} = R \left(\frac{1}{n^2} - \frac{1}{(n + \delta n)^2} \right); \quad \delta n = \delta r / r_0$	<p>Unknown</p>
18	<p>The nature of the Lamb shift: the shift is precisely equal to the frequency gaps between the nearest spectral terms of the background spectrum (see # 17)</p>	<p>An erroneous concept based on an influence of the invented (non-existed) virtual particles</p>
19	<p>The precise derivation of binding energy in atoms without use of the relation</p> $\Delta E = \Delta m \cdot c^2$	<p>Unable</p>
20	<p>The physical meaning of the speed of light c in the relation</p> $E_0 = m_0 c^2;$ <p>m_0 is the associated mass of a particle (see # 1). Speed of light c is the basis wave speed of exchange of matter-space-time at the subatomic level.</p>	<p>Unknown</p> <p>m_0 is the “rest” mass.</p>

21	<p>Internal spatial structure of atoms, <i>i. e.</i>, the disposition of nucleons in atoms (The latter defines the structural variety at the molecular level in Nature: “genetic code”)</p>	<p>The fixed (strictly geometrical) disposition of nucleons is Unknown</p>
22	<p>The g-lepton structure of nucleons: Proton and Neutron are similar in g-lepton structure to isotopes $^{28}_{14}\text{Si}$ and $^{29}_{14}\text{Si}$, respectively, according to Shell-Nodal Atomic Model $(m_g = 68.22 m_e)$</p>	<p>Quark structure (does not similar to crystal)</p>
23	<p>Crystal structure of solids, including forbidden by mathematical laws of crystallography</p>	<p>Unable</p>
24	<p>The structure of all isotopes and their relative masses (including limiting masses: minimal and maximal for every isotope)</p>	<p>Unable</p>
25	<p>The nature of Mendeleev’s Periodic Law: the similarity of nodal structure of external atomic nucleon shells.</p>	<p>Different explanation: electron structure of atoms</p>
26	<p>The fine structure constant physical meaning: the scale correlation between basis and superstructure of wave (between oscillatory and wave processes in waves)</p>	<p>Unknown</p>
27	<p>The unified description of electromagnetic, gravitational, and strong (nuclear) interactions</p>	<p>Unable</p>

28	<p><i>The nature of the spherical harmonics of wave and Schrodinger equations</i></p> <p>The spherical harmonics define polar-azimuthal coordinates of nodes and antinodes of standing spherical waves</p>	<p><i>Unknown</i></p> <p>As a result, an introduction in quantum mechanics of the conceptually unfounded notion of hybridization of atomic orbitals</p>
29	<p><i>The nature of integer and fractional quantization in quantum Hall effect</i></p> <p>The nature of quantization in the Hall conductance (the resistance quantum) is naturally uncovered as an internal feature of atomic structures considered as wave formations, without accounting an influence of external magnetic fields.</p> <p><i>The deduced spectrum of fundamental resistances</i></p> $R_e = \frac{h}{e^2} \frac{m}{n}$	<p><i>Fitting theory in the spirit of the virtual particles of quantum electrodynamics</i></p> <p>Modern explanation is based on an imaginary quantum-mechanical fluid of a hypothetical new form and on a many body wave function. It predicts that the elementary excitations involve pseudo-particle charge carriers with charges that are fractions of the electronic charge.</p>
30	<p><i>Precise derivation of the neutron magnetic moment</i></p> $\mu_n(th) = \frac{e\nu_0}{c} \left[\left(\tilde{\lambda}_e + \frac{r_0}{y_{0,12}} \right) \sqrt{\frac{2Rh}{m_0c}} + \frac{r_e}{j_{0,12}} \sqrt{\frac{2Rh_e}{m_0c}} \right]$ $\mu_n(th) = -0.96623513 \cdot 10^{-26} J \cdot T^{-1}$	<p><i>Unable</i></p>
31	<p><i>Precise derivation of the proton magnetic moment</i></p> $\mu_p(th) = \frac{(e + \Delta e_p)\nu_0}{c} \left(\tilde{\lambda}_e + r_0 \frac{1}{\beta} \frac{(a'_{0,11} + y_{0,12})}{2(a'_{0,11}y_{0,12})} \right) \sqrt{\frac{2Rh}{m_0c}}$ $\mu_p(th) = 1.410606662 \cdot 10^{-26} J \cdot T^{-1}$	<p><i>Unable</i></p>

32 **Objective (true) dimensionalities of physical quantities in integer powers of units of matter (g), space (cm), and time (s):**

Electric charge, $[q] = [m]/[t] = g \cdot s^{-1}$

Electric current, $[I] = [q]/[t] = g \cdot s^{-2}$

Circulation, $[\Gamma] = [I]/[c] = g \cdot cm^{-1} \cdot s^{-1}$

Electric field strength, $[E] = [F]/[q] = cm \cdot s^{-1}$

Magnetic field strength, $[B] = [F]/[q] = cm \cdot s^{-1}$

Electric field momentum density, $[D] = [\epsilon_0][E] = g \cdot cm^{-2} \cdot s^{-1}$

Magnetic field momentum density, $[H] = [\epsilon_0][B] = g \cdot cm^{-2} \cdot s^{-1}$

Potential, $[U] = [F][l]/[q] = cm^2 \cdot s^{-1}$

Resistance, $[R] = [U]/[I] = g^{-1} \cdot cm^2 \cdot s$

Conductance, $[G] = [R]^{-1} = g \cdot cm^{-2} \cdot s^{-1}$

Resistivity, $[\rho] = [R][l] = g^{-1} \cdot cm^3 \cdot s$

Conductivity, $[\sigma] = [\rho]^{-1} = g \cdot cm^{-3} \cdot s^{-1}$

Inductance, $[L] = [U][t]/[I] = g^{-1} \cdot cm^2 \cdot s^2$

etc.

Other physical quantities of electromagnetism contained electric charge, current, and their derivatives with corrected dimensionalities.

Incorrect dimensionalities (subjective, phenomenological)

Accepted in contemporary physics dimensionalities of physical quantities of electromagnetism, based on the erroneous dimensionalities of electric charge, current, and their derivatives, are erroneous.

33 **Fundamental Period of the Decimal Code of the Universe**

$$\Delta = 2\pi \lg e = 2.7287527\dots$$

Unknown

Literature

- [1] L. Kreidik and G. Shpenkov, *Dynamic Model of Elementary Particles and the Nature of Mass and 'Electric' Charge*, "Revista Ciências Exatas e Naturais", Vol. 3, No 2, 157-170, (2001); <http://www.unicentro.br/editora/revistas/recen/v3n2/trc510final.pdf>
- [2] L. G. Kreidik and G. P. Shpenkov, *Atomic Structure of Matter-Space*, Geo. S., Bydgoszcz, 2001, 584 p.
- [3] G. P. Shpenkov and L. G. Kreidik, *On Electron Spin of $\hbar/2$* , Hadronic Journal, Vol. 25, No. 5, 573-586, (2002).
- [4] G. P. Shpenkov, *Shell-Nodal Atomic Model*, Hadronic Journal Supplement, **17**, No. 4, 507-566 (2002).
- [5] L. G. Kreidik and G. P. Shpenkov, *Microwave Background Radiation of Hydrogen Atoms*, Revista Ciências Exatas e Naturais, **4**, No. 1, 9-18 (2002); www.unicentro.br/pesquisa/editora/revistas/exatas/v4n1/Microwave.pdf
- [6] L. G. Kreidik and G. P. Shpenkov, *Philosophy of Contents-Form and Coulomb's Law*, Proceedings of The Twentieth World Congress of Philosophy, Copley Place, Boston, Massachusetts, USA, 10-16 August, 1998; <http://www.bu.edu/wcp/Papers/Scie/ScieShpe.htm>
- [7] G. P. Shpenkov, *Derivation of the Lamb Shift with Due Account of Wave Features for the Proton-Electron Interaction*; Revista Ciências Exatas e Naturais, Vol. 6, No 2, 171-185, (2004); <http://shpenkov.janmax.com/derivation.pdf>
- [8] G. P. Shpenkov, *An Elucidation of the Nature of the Periodic Law*, Chapter 7 in "*The Mathematics of the Periodic Table*", edited by Rouvray D. H. and King R. B., Nova Science Publishers, NY, pp. 119-160, 2006.
- [9] G. P. Shpenkov, *The first Precise Derivation of the Magnetic Moment of an Electron Beyond Quantum Electrodynamics*, Physics Essays, **19**, No. 1, (2006).
- [10] G. P. Shpenkov, *On the Fine-Structure Constant Physical Meaning*, Hadronic Journal, Vol. 28, No. 3, 337-372, (2005).
- [11] G. P. Shpenkov and L. G. Kreidik, *Conjugate Parameters of Physical Processes and Physical Time*, Physics Essays, **15**, No. 3, 339-349, (2002).
- [12] L. G. Kreidik and G. P. Shpenkov, *Important Results of Analyzing Foundations of Quantum Mechanics*, Galilean Electrodynamics & QED-EAST, Vol. 13, SI No. 2, 23-30, (2002); <http://shpenkov.janmax.com/QM-Analysis.pdf>
- [13] G. P. Shpenkov and L. G. Kreidik, *Schrödinger's Errors of Principle*, Galilean Electrodynamics, Vol. 16, No. 3, 51-56, (2005); <http://shpenkov.janmax.com/Blunders.pdf>
- [14] G. P. Shpenkov and L. G. Kreidik, *What the Electric Charge is*; <http://shpenkov.janmax.com/Elec-Charge.pdf>
- [15] G. P. Shpenkov, *Conceptual Unfoundedness of Hybridization and the Nature of the Spherical Harmonics*, HADRONIC JOURNAL, Vol. 29. No. 4, p. 455, (2006).
- [16] G. P. Shpenkov, *The Dependence of Hall Conductance Quanta on the Fundamental Frequency of the Atomic Level*, (2009); <http://shpenkov.janmax.com/Hall.pdf>
- [17] G. P. Shpenkov, *Derivation of Neutron's Magnetic Moment on the Basis of Dynamic Model of Elementary Particles*, (2008); <http://shpenkov.janmax.com/neutronmagnmom.pdf>
- [18] G. P. Shpenkov, *Derivation of the Proton's Magnetic Moment beyond QED and QCD Theories*, (2008); <http://shpenkov.janmax.com/protonmagnmom.pdf>

See also: <http://shpenkov.janmax.com/selectedpub.asp>